

Small molecule modulators of RING-type E3 ligases: MDM and cullin families as targets

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2018 Bulatov, Zagidullin, Valiullina, Sayarova and Rizvanov. Ubiquitin-proteasome system (UPS) is a primary signaling pathway for regulation of intracellular protein levels. E3 ubiquitin ligases, substrate-specific members of the UPS, represent highly attractive protein targets for drug discovery. The importance of E3 ligases as prospective targets for small molecule modulation is reinforced by ever growing evidence of their role in cancer and other diseases. To date the number of potent compounds targeting E3 ligases remains rather low and their rational design constitutes a challenging task. To successfully address this problem one must take into consideration the multi-subunit nature of many E3 ligases that implies multiple druggable pockets and protein-protein interfaces. In this review, we briefly cover the current state of drug discovery in the field of RING-type E3 ligases with focus on MDM and Cullin families as targets. We also provide an overview of small molecule chimeras that induce RING-type E3-mediated proteasomal degradation of substrate proteins of interest.

<http://dx.doi.org/10.3389/fphar.2018.00450>

Keywords

Cullin family, Induced protein degradation, MDM family, PROTACs, RING-type E3 ligases, Small molecules, SNIPERs, Ubiquitin-proteasome system

References

- [1] Andérica-Romero, A. C., González-Herrera, I. G., Santamaría, A., and Pedraza-Chaverri, J. (2013). Cullin 3 as a novel target in diverse pathologies. *Redox Biol.* 1, 366-372. doi: 10.1016/j.redox.2013.07.003
- [2] Bernal, F., Wade, M., Godes, M., Davis, T. N., Whitehead, D. G., Kung, A. L., et al. (2010). A stapled p53 helix overcomes HDMX-mediated suppression of p53. *Cancer Cell* 18, 411-422. doi: 10.1016/j.ccr.2010.10.024
- [3] Bista, M., Smithson, D., Pecak, A., Salinas, G., Pustelny, K., Min, J., et al. (2012). On the mechanism of action of SJ-172550 in inhibiting the interaction of MDM4 and p53. *PLoS One* 7:e37518. doi: 10.1371/journal.pone.0037518
- [4] Bondeson, D. P., Mares, A., Smith, I. E. D., Ko, E., Campos, S., Miah, A. H., et al. (2015). Catalytic in vivo protein knockdown by small-molecule PROTACs. *Nat. Chem. Biol.* 11, 611-617. doi: 10.1038/nchembio.1858
- [5] Bosu, D. R., and Kipreos, E. T. (2008). Cullin-RING ubiquitin ligases: global regulation and activation cycles. *Cell Div.* 3:7. doi: 10.1186/1747-102-3-7
- [6] Buckley, D. L., and Crews, C. M. (2014). Small-molecule control of intracellular protein levels through modulation of the ubiquitin proteasome system. *Angew. Chem. Int. Ed. Engl.* 53, 2312-2330. doi: 10.1002/anie.201307761
- [7] Buckley, D. L., Van Molle, I., Gareiss, P. C., Tae, H. S., Michel, J., Noblin, D. J., et al. (2012). Targeting the von Hippel-Lindau E3 ubiquitin ligase using small molecules to disrupt the VHL/HIF-1 α interaction. *J. Am. Chem. Soc.* 134, 4465-4468. doi: 10.1021/ja209924v
- [8] Bulatov, E., and Ciulli, A. (2015). Targeting Cullin-RING E3 ubiquitin ligases for drug discovery: structure, assembly and small-molecule modulation. *Biochem. J.* 467, 365-386. doi: 10.1042/BJ20141450
- [9] Bulatov, E., Martin, E. M., Chatterjee, S., Knebel, A., Shimamura, S., Konijnenberg, A., et al. (2015). Biophysical studies on interactions and assembly of full-size E3 ubiquitin ligase: suppressor of cytokine signaling 2 (SOCS2)-elongin BC-cullin 5-ring box protein 2 (RBX2). *J. Biol. Chem.* 290, 4178-4191. doi: 10.1074/jbc.M114.616664
- [10] Cardote, T. A. F., Gadd, M. S., and Ciulli, A. (2017). Crystal structure of the Cul2-Rbx1-EloBC-VHL ubiquitin ligase complex. *Structure* 25, 901.e3-911.e3. doi: 10.1016/j.str.2017.04.009

- [11] Cardozo, T., and Abagyan, R. (2005). Druggability of SCF ubiquitin ligase-protein interfaces. *Methods Enzymol.* 399, 634-653. doi: 10.1016/S0076-6879(05)99042-3
- [12] Castro, A., Bernis, C., Vigneron, S., Labbé, J.-C., and Lorca, T. (2005). The anaphase-promoting complex: a key factor in the regulation of cell cycle. *Oncogene* 24, 314-325. doi: 10.1038/sj.onc.1207973
- [13] Chamberlain, P. P., Lopez-Girona, A., Miller, K., Carmel, G., Pagarigan, B., Chie-Leon, B., et al. (2014). Structure of the human Cereblon-DDB1-lenalidomide complex reveals basis for responsiveness to thalidomide analogs. *Nat. Struct. Mol. Biol.* 21, 803-809. doi: 10.1038/nsmb.2874
- [14] Chan, K.-H., Zengerle, M., Testa, A., and Ciulli, A. (2018). Impact of target warhead and linkage vector on inducing protein degradation: comparison of bromodomain and extra-terminal (BET) degraders derived from triazolodiazepine (JQ1) and tetrahydroquinoline (I-BET726) BET inhibitor scaffolds. *J. Med. Chem.* 61, 504-513. doi: 10.1021/acs.jmedchem.6b01912
- [15] Chang, Y. S., Graves, B., Guerlavais, V., Tovar, C., Packman, K., To, K.-H., et al. (2013). Stapled α -helical peptide drug development: a potent dual inhibitor of MDM2 and MDMX for p53-dependent cancer therapy. *Proc. Natl. Acad. Sci. U.S.A.* 110, E3445-E3454. doi: 10.1073/pnas.1303002110
- [16] Chen, Q., Xie, W., Kuhn, D. J., Voorhees, P. M., Lopez-Girona, A., Mendy, D., et al. (2008). Targeting the p27 E3 ligase SCF(Skp2) results in p27- and Skp2-mediated cell-cycle arrest and activation of autophagy. *Blood* 111, 4690-4699. doi: 10.1182/blood-2007-09-112904
- [17] Cohen, P., and Tcherpakov, M. (2010). Will the ubiquitin system furnish as many drug targets as protein kinases? *Cell* 143, 686-693. doi: 10.1016/j.cell.2010.11.016
- [18] Crews, C. M. (2018). Inducing protein degradation as a therapeutic strategy. *J. Med. Chem.* 61, 403-404. doi: 10.1021/acs.jmedchem.7b01333
- [19] Deshaies, R. J., and Joazeiro, C. A. P. (2009). RING domain E3 ubiquitin ligases. *Annu. Rev. Biochem.* 78, 399-434. doi: 10.1146/annurev.biochem.78.101807.093809
- [20] Duda, D. M., Borg, L. A., Scott, D. C., Hunt, H. W., Hammel, M., and Schulman, B. A. (2008). Structural insights into NEDD8 activation of cullin-RING ligases: conformational control of conjugation. *Cell* 134, 995-1006. doi: 10.1016/j.cell.2008.07.022
- [21] Duda, D. M., Scott, D. C., Calabrese, M. F., Zimmerman, E. S., Zheng, N., and Schulman, B. A. (2011). Structural regulation of cullin-RING ubiquitin ligase complexes. *Curr. Opin. Struct. Biol.* 21, 257-264. doi: 10.1016/j.sbi.2011.01.003
- [22] Feldman, R. M., Correll, C. C., Kaplan, K. B., and Deshaies, R. J. (1997). A complex of Cdc4p, Skp1p, and Cdc53p/cullin catalyzes ubiquitination of the phosphorylated CDK inhibitor Sic1p. *Cell* 91, 221-230. doi: 10.1016/S0092-8674(00)80404-3
- [23] Fischer, E. S., Böhm, K., Lydeard, J. R., Yang, H., Stadler, M. B., Cavadini, S., et al. (2014). Structure of the DDB1-CRBN E3 ubiquitin ligase in complex with thalidomide. *Nature* 512, 49-53. doi: 10.1038/nature13527
- [24] Fischer, E. S., Scrima, A., Böhm, K., Matsumoto, S., Lingaraju, G. M., Faty, M., et al. (2011). The molecular basis of CRL4DDB2/CSA ubiquitin ligase architecture, targeting, and activation. *Cell* 147, 1024-1039. doi: 10.1016/j.cell.2011.10.035
- [25] Frost, J., Galdeano, C., Soares, P., Gadd, M. S., Grzes, K. M., Ellis, L., et al. (2016). Potent and selective chemical probe of hypoxic signalling downstream of HIF- α hydroxylation via VHL inhibition. *Nat. Commun.* 7:13312. doi: 10.1038/ncomms13312
- [26] Gadd, M. S., Bulatov, E., and Ciulli, A. (2015). Serendipitous SAD Solution for DMSO-Soaked SOCS2-ElonginC-ElonginB crystals using covalently incorporated dimethylarsenic: insights into substrate receptor conformational flexibility in cullin RING ligases. *PLoS One* 10:e0131218. doi: 10.1371/journal.pone.0131218
- [27] Gadd, M. S., Testa, A., Lucas, X., Chan, K.-H., Chen, W., Lamont, D. J., et al. (2017). Structural basis of PROTAC cooperative recognition for selective protein degradation. *Nat. Chem. Biol.* 13, 514-521. doi: 10.1038/nchembio.2329
- [28] Galdeano, C., Gadd, M. S., Soares, P., Scaffidi, S., Van Molle, I., Birced, I., et al. (2014). Structure-guided design and optimization of small molecules targeting the protein-protein interaction between the von Hippel-Lindau (VHL) E3 ubiquitin ligase and the hypoxia inducible factor (HIF) α subunit with in vitro nanomolar affinities. *J. Med. Chem.* 57, 8657-8663. doi: 10.1021/jm5011258
- [29] Geyer, R. K., Yu, Z. K., and Maki, C. G. (2000). The MDM2 RING-finger domain is required to promote p53 nuclear export. *Nat. Cell Biol.* 2, 569-573. doi: 10.1038/35023507
- [30] Hardcastle, I. R., Liu, J., Valeur, E., Watson, A., Ahmed, S. U., Blackburn, T. J., et al. (2011). Isoindolinone inhibitors of the murine double minute 2 (MDM2)-p53 protein-protein interaction: structure-activity studies leading to improved potency. *J. Med. Chem.* 54, 1233-1243. doi: 10.1021/jm1011929
- [31] Hatakeyama, S., Yada, M., Matsumoto, M., Ishida, N., and Nakayama, K. I. (2001). U box proteins as a new family of ubiquitin-protein ligases. *J. Biol. Chem.* 276, 33111-33120. doi: 10.1074/jbc.M102755200
- [32] Haupt, S., Vijayakumaran, R., Miranda, P. J., Burgess, A., Lim, E., and Haupt, Y. (2017). The role of MDM2 and MDM4 in breast cancer development and prevention. *J. Mol. Cell Biol.* 9, 53-61. doi: 10.1093/jmcb/mjx007
- [33] Hayashi, K.-I., Tan, X., Zheng, N., Hatate, T., Kimura, Y., Kepinski, S., et al. (2008). Small-molecule agonists and antagonists of F-box protein-substrate interactions in auxin perception and signaling. *Proc. Natl. Acad. Sci. U.S.A.* 105, 5632-5637. doi: 10.1073/pnas.0711146105
- [34] Hershko, A., and Ciechanover, A. (1998). The ubiquitin system. *Annu. Rev. Biochem.* 67, 425-479. doi: 10.1146/annurev.biochem.67.1.425
- [35] Holzer, P., Masuya, K., Furet, P., Kallen, J., Valat-Stachyra, T., Ferretti, S., et al. (2015). Discovery of a dihydroisoquinolinone derivative (NVP-CGM097): a highly potent and selective MDM2 inhibitor undergoing phase 1 clinical trials in p53wt tumors. *J. Med. Chem.* 58, 6348-6358. doi: 10.1021/acs.jmedchem.5b00810
- [36] Hu, B., Gilkes, D. M., and Chen, J. (2007). Efficient p53 activation and apoptosis by simultaneous disruption of binding to MDM2 and MDMX. *Cancer Res.* 67, 8810-8817. doi: 10.1158/0008-5472.CAN-07-1140
- [37] Huang, L., Yan, Z., Liao, X., Li, Y., Yang, J., Wang, Z.-G., et al. (2011). The p53 inhibitors MDM2/MDMX complex is required for control of p53 activity in vivo. *Proc. Natl. Acad. Sci. U.S.A.* 108, 12001-12006. doi: 10.1073/pnas.1102309108
- [38] Itoh, Y., Ishikawa, M., Naito, M., and Hashimoto, Y. (2010). Protein knockdown using methyl bestatin-ligand hybrid molecules: design and synthesis of inducers of ubiquitination-mediated degradation of cellular retinoic acid-binding proteins. *J. Am. Chem. Soc.* 132, 5820-5826. doi: 10.1021/ja100691p
- [39] Itoh, Y., Kitaguchi, R., Ishikawa, M., Naito, M., and Hashimoto, Y. (2011). Design, synthesis and biological evaluation of nuclear receptor-degradation inducers. *Bioorg. Med. Chem.* 19, 6768-6778. doi: 10.1016/j.bmc.2011.09.041
- [40] Kamura, T., Koeppe, D. M., Conrad, M. N., Skowry, D., Moreland, R. J., Iliopoulos, O., et al. (1999). Rbx1, a component of the VHL tumor suppressor complex and SCF ubiquitin ligase. *Science* 284, 657-661. doi: 10.1126/science.284.5414.657
- [41] Lai, A. C., and Crews, C. M. (2017). Induced protein degradation: an emerging drug discovery paradigm. *Nat. Rev. Drug Discov.* 16, 101-114. doi: 10.1038/nrd.2016.211
- [42] Lai, A. C., Toure, M., Hellerschmied, D., Salami, J., Jaime-Figueroa, S., Ko, E., et al. (2016). Modular PROTAC design for the degradation of oncogenic BCR-ABL. *Angew. Chem. Int. Ed. Engl.* 55, 807-810. doi: 10.1002/anie.201507634
- [43] Lee, G. M., and Craik, C. S. (2009). Trapping moving targets with small molecules. *Science* 324, 213-215. doi: 10.1126/science.1169378
- [44] Li, Z., and Xiong, Y. (2017). Cytoplasmic E3 ubiquitin ligase CUL9 controls cell proliferation, senescence, apoptosis and genome integrity through p53. *Oncogene* 36, 5212-5218. doi: 10.1038/onc.2017.141
- [45] Lu, J., Qian, Y., Altieri, M., Dong, H., Wang, J., Raina, K., et al. (2015). Hijacking the E3 ubiquitin ligase cereblon to efficiently target BRD4. *Chem. Biol.* 22, 755-763. doi: 10.1016/j.chembiol.2015.05.009
- [46] Lydeard, J. R., Schulman, B. A., and Harper, J. W. (2013). Building and remodelling Cullin-RING E3 ubiquitin ligases. *EMBO Rep.* 14, 1050-1061. doi: 10.1038/embor.2013.173

- [47] Maniaci, C., Hughes, S. J., Testa, A., Chen, W., Lamont, D. J., Rocha, S., et al. (2017). Homo-PROTACs: bivalent small-molecule dimerizers of the VHL E3 ubiquitin ligase to induce self-degradation. *Nat. Commun.* 8:830. doi: 10.1038/s41467-017-00954-1
- [48] Metzger, M. B., Pruneda, J. N., Klevit, R. E., and Weissman, A. M. (2014). RING-type E3 ligases: master manipulators of E2 ubiquitin-conjugating enzymes and ubiquitination. *Biochim. Biophys. Acta* 1843, 47-60. doi: 10.1016/j.bbamcr.2013.05.026
- [49] Muller, P. A. J., and Vousden, K. H. (2013). p53 mutations in cancer. *Nat. Cell Biol.* 15, 2-8. doi: 10.1038/ncb2641
- [50] Ohoka, N., Misawa, T., Kurihara, M., Demizu, Y., and Naito, M. (2017a). Development of a peptide-based inducer of protein degradation targeting NOTCH1. *Bioorg. Med. Chem. Lett.* 27, 4985-4988. doi: 10.1016/j.bmcl.2017.10.011
- [51] Ohoka, N., Okuhira, K., Ito, M., Nagai, K., Shibata, N., Hattori, T., et al. (2017b). In vivo knockdown of pathogenic proteins via specific and nongenetic inhibitor of apoptosis protein (IAP)-dependent protein erasers (SNIPERs). *J. Biol. Chem.* 292, 4556-4570. doi: 10.1074/jbc.M116.768853
- [52] Ohoka, N., Morita, Y., Nagai, K., Shimokawa, K., Ujikawa, O., Fujimori, I., et al. (2018). Derivatization of inhibitor of apoptosis protein (IAP) ligands yields improved inducers of estrogen receptor degradation. *J. Biol. Chem.* doi: 10.1074/jbc.RA117.001091 [Epub ahead of print]
- [53] Ohta, T., Michel, J. J., Schottelius, A. J., and Xiong, Y. (1999). ROC1, a homolog of APC11, represents a family of cullin partners with an associated ubiquitin ligase activity. *Mol. Cell* 3, 535-541. doi: 10.1016/S1097-2765(00)80482-7
- [54] Okuhira, K., Ohoka, N., Sai, K., Nishimaki-Mogami, T., Itoh, Y., Ishikawa, M., et al. (2011). Specific degradation of CRABP-II via cIAP1-mediated ubiquitylation induced by hybrid molecules that crosslink cIAP1 and the target protein. *FEBS Lett.* 585, 1147-1152. doi: 10.1016/j.febslet.2011.03.019
- [55] Pant, V., Xiong, S., Iwakuma, T., Quintás-Cardama, A., and Lozano, G. (2011). Heterodimerization of Mdm2 and Mdm4 is critical for regulating p53 activity during embryogenesis but dispensable for p53 and Mdm2 stability. *Proc. Natl. Acad. Sci. U.S.A.* 108, 11995-12000. doi: 10.1073/pnas.1102241108
- [56] Pellegrino, M., Mancini, F., Lucà, R., Coletti, A., Giacchè, N., Manni, I., et al. (2015). Targeting the MDM2/MDM4 interaction interface as a promising approach for p53 reactivation therapy. *Cancer Res.* 75, 4560-4572. doi: 10.1158/0008-5472.CAN-15-0439
- [57] Petroski, M. D. (2008). The ubiquitin system, disease, and drug discovery. *BMC Biochem.* 9(Suppl. 1):S7. doi: 10.1186/1471-2091-9-S1-S7
- [58] Pickart, C. M. (2001). Mechanisms underlying ubiquitination. *Annu. Rev. Biochem.* 70, 503-533. doi: 10.1146/annurev.biochem.70.1.503
- [59] Popowicz, G. M., Czarna, A., Wolf, S., Wang, K., Wang, W., Dömling, A., et al. (2010). Structures of low molecular weight inhibitors bound to MDMX and MDM2 reveal new approaches for p53-MDMX/MDM2 antagonist drug discovery. *Cell Cycle* 9, 1104-1111. doi: 10.4161/cc.9.6.10956
- [60] Raina, K., Lu, J., Qian, Y., Altieri, M., Gordon, D., Rossi, A. M. K., et al. (2016). PROTAC-induced BET protein degradation as a therapy for castration-resistant prostate cancer. *Proc. Natl. Acad. Sci. U.S.A.* 113, 7124-7129. doi: 10.1073/pnas.1521738113
- [61] Reed, D., Shen, Y., Shelat, A. A., Arnold, L. A., Ferreira, A. M., Zhu, F., et al. (2010). Identification and characterization of the first small molecule inhibitor of MDMX. *J. Biol. Chem.* 285, 10786-10796. doi: 10.1074/jbc.M109.056747
- [62] Sarikas, A., Hartmann, T., and Pan, Z.-Q. (2011). The cullin protein family. *Genome Biol.* 12:220. doi: 10.1186/gb-2011-12-4-220
- [63] Schneekloth, A. R., Puchault, M., Tae, H. S., and Crews, C. M. (2008). Targeted intracellular protein degradation induced by a small molecule: en route to chemical proteomics. *Bioorg. Med. Chem. Lett.* 18, 5904-5908. doi: 10.1016/j.bmcl.2008.07.114
- [64] Seol, J. H., Feldman, R. M., Zachariae, W., Shevchenko, A., Correll, C. C., Lyapina, S., et al. (1999). Cdc53/cullin and the essential Hrt1 RING-H2 subunit of SCF define a ubiquitin ligase module that activates the E2 enzyme Cdc34. *Genes Dev.* 13, 1614-1626. doi: 10.1101/gad.13.12.1614
- [65] Shibata, N., Miyamoto, N., Nagai, K., Shimokawa, K., Sameshima, T., Ohoka, N., et al. (2017). Development of protein degradation inducers of oncogenic BCR-ABL protein by conjugation of ABL kinase inhibitors and IAP ligands. *Cancer Sci.* 108, 1657-1666. doi: 10.1111/cas.13284
- [66] Shibata, N., Nagai, K., Morita, Y., Ujikawa, O., Ohoka, N., Hattori, T., et al. (2018). Development of protein degradation inducers of androgen receptor by conjugation of androgen receptor ligands and inhibitor of apoptosis protein ligands. *J. Med. Chem.* 61, 543-575. doi: 10.1021/acs.jmedchem.7b00168
- [67] Soares, P., Gadd, M. S., Frost, J., Galdeano, C., Ellis, L., Epemolu, O., et al. (2018). Group-based optimization of potent and cell-active inhibitors of the von Hippel-Lindau (VHL) E3 ubiquitin ligase: structure-activity relationships leading to the chemical probe (2S,4R)-1-((S)--(1-cyanocyclopropanecarboxamido)-3,3-dimethylbutanoyl)-4-hydroxy-N-(4-(4-methylthiazol-5-yl)benzyl)pyrrolidine-2-carboxamide (VH298). *J. Med. Chem.* 61, 599-618. doi: 10.1021/acs.jmedchem.7b00675
- [68] Soucy, T. A., Smith, P. G., Milhollen, M. A., Berger, A. J., Gavin, J. M., Adhikari, S., et al. (2009). An inhibitor of NEDD8-activating enzyme as a new approach to treat cancer. *Nature* 458, 732-736. doi: 10.1038/nature07884
- [69] Spratt, D. E., Walden, H., and Shaw, G. S. (2014). RBR E3 ubiquitin ligases: new structures, new insights, new questions. *Biochem. J.* 458, 421-437. doi: 10.1042/BJ20140006
- [70] Sun, Y. (2005). Overview of approaches for screening for ubiquitin ligase inhibitors. *Methods Enzymol.* 399, 654-663. doi: 10.1016/S0076-6879(05)99043-5
- [71] Swords, R. T., Erba, H. P., DeAngelo, D. J., Bixby, D. L., Altman, J. K., Maris, M., et al. (2015). Pevonedistat (MLN4924), a First-in-Class NEDD8-activating enzyme inhibitor, in patients with acute myeloid leukaemia and myelodysplastic syndromes: a phase 1 study. *Br. J. Haematol.* 169, 534-543. doi: 10.1111/bjh.13323
- [72] Tan, P., Fuchs, S. Y., Chen, A., Wu, K., Gomez, C., Ronai, Z., et al. (1999). Recruitment of a ROC1-CUL1 ubiquitin ligase by Skp1 and HOS to catalyze the ubiquitination of I κ B α . *Mol. Cell* 3, 527-533. doi: 10.1016/S1097-2765(00)80481-5
- [73] Thiel, P., Kaiser, M., and Ottmann, C. (2012). Small-molecule stabilization of protein-protein interactions: an underestimated concept in drug discovery? *Angew. Chem. Int. Ed. Engl.* 51, 2012-2018. doi: 10.1002/anie.201107616
- [74] Tisato, V., Voltan, R., Gonelli, A., Secchiero, P., and Zauli, G. (2017). MDM2/X inhibitors under clinical evaluation: perspectives for the management of hematological malignancies and pediatric cancer. *J. Hematol. Oncol.* 10:133. doi: 10.1186/s13045-017-0500-5
- [75] Van Molle, I., Thomann, A., Buckley, D. L., So, E. C., Lang, S., Crews, C. M., et al. (2012). Dissecting fragment-based lead discovery at the von Hippel-Lindau protein:hypoxia inducible factor 1 α protein-protein interface. *Chem. Biol.* 19, 1300-1312. doi: 10.1016/j.chembiol.2012.08.015
- [76] Vassilev, L. T., Vu, B. T., Graves, B., Carvajal, D., Podlaski, F., Filipovic, Z., et al. (2004). In vivo activation of the p53 pathway by small-molecule antagonists of MDM2. *Science* 303, 844-848. doi: 10.1126/science.1092472
- [77] Vu, B., Wovkulich, P., Pizzolato, G., Lovey, A., Ding, Q., Jiang, N., et al. (2013). Discovery of RG7112: a small-molecule MDM2 inhibitor in clinical development. *ACS Med. Chem. Lett.* 4, 466-469. doi: 10.1021/ml4000657
- [78] Winter, G. E., Buckley, D. L., Paulk, J., Roberts, J. M., Souza, A., Dhe-Paganon, S., et al. (2015). DRUG DEVELOPMENT. Phthalimide conjugation as a strategy for in vivo target protein degradation. *Science* 348, 1376-1381. doi: 10.1126/science.aab1433
- [79] Xirodimas, D. P., Saville, M. K., Bourdon, J.-C., Hay, R. T., and Lane, D. P. (2004). Mdm2-mediated NEDD8 conjugation of p53 inhibits its transcriptional activity. *Cell* 118, 83-97. doi: 10.1016/j.cell.2004.06.016
- [80] Zaytsev, A., Dodd, B., Magnani, M., Ghiron, C., Golding, B. T., Griffin, R. J., et al. (2015). Searching for dual inhibitors of the MDM2-p53 and MDMX-p53 protein-protein interaction by a scaffold-hopping approach. *Chem. Biol. Drug Des.* 86, 180-189. doi: 10.1111/cbdd.12474
- [81] Zengerle, M., Chan, K.-H., and Ciulli, A. (2015). Selective small molecule induced degradation of the BET bromodomain protein BRD4. *ACS Chem. Biol.* 10, 1770-1777. doi: 10.1021/acschembio.5b00216
- [82] Zhang, B., Golding, B. T., and Hardcastle, I. R. (2015). Small-molecule MDM2-p53 inhibitors: recent advances. *Future Med. Chem.* 7, 631-645. doi: 10.4155/fmc.15.13
- [83] Zhao, Y., Liu, L., Sun, W., Lu, J., McEachern, D., Li, X., et al. (2013). Diastereomeric spirooxindoles as highly potent and efficacious MDM2 inhibitors. *J. Am. Chem. Soc.* 135, 7223-7234. doi: 10.1021/ja3125417

- [84] Zhao, Y., and Sun, Y. (2013). Cullin-RING Ligases as attractive anti-cancer targets. *Curr. Pharm. Des.* 19, 3215-3225. doi: 10.2174/13816128113199990300
- [85] Zheng, N., Schulman, B. A., Song, L., Miller, J. J., Jeffrey, P. D., Wang, P., et al. (2002). Structure of the Cul1-Rbx1-Skp1-F boxSkp2 SCF ubiquitin ligase complex. *Nature* 416, 703-709. doi: 10.1038/416703a
- [86] Zhou, B., Hu, J., Xu, F., Chen, Z., Bai, L., Fernandez-Salas, E., et al. (2017). Discovery of a small-molecule degrader of bromodomain and extra-terminal (BET) proteins with picomolar cellular potencies and capable of achieving tumor regression. *J. Med. Chem.* 61, 462-481. doi: 10.1021/acs.jmedchem.6b01816